

Differentiation-induced replication-timing changes are restricted to AT-rich/long interspersed nuclear element (LINE)-rich isochores

Ichiro Hiratani, Amanda Leskovar, and David M. Gilbert*

Department of Biochemistry and Molecular Biology, State University of New York Upstate Medical University, 750 East Adams Street, Syracuse, NY 13210

Edited by Mark T. Groudine, Fred Hutchinson Cancer Research Center, Seattle, WA, and approved October 20, 2004 (received for review September 9, 2004)

The replication timing of some genes is developmentally regulated, but the significance of replication timing to cellular differentiation has been difficult to substantiate. Studies have largely been restricted to the comparison of a few genes in established cell lines derived from different tissues, and most of these genes do not change replication timing. Hence, it has not been possible to predict how many or what types of genes might be subject to such control. Here, we have evaluated the replication timing of 54 tissue-specific genes in mouse embryonic stem cells before and after differentiation to neural precursors. Strikingly, genes residing within isochores rich in GC and poor in long interspersed nuclear elements (LINEs) did not change their replication timing, whereas half of genes within isochores rich in AT and long interspersed nuclear elements displayed programmed changes in replication timing that accompanied changes in gene expression. Our results provide direct evidence that differentiation-induced autosomal replication-timing changes are a significant part of mammalian development, provide a means to predict genes subject to such regulation, and suggest that replication timing may be more related to the evolution of metazoan genomes than to gene function or expression pattern.

It is generally presumed that early replication is a necessary (albeit not sufficient) condition for transcription, whereas late replicating sequences are assembled into transcriptionally inactive chromatin; however, the evidence for this assumption has been far from conclusive (1, 2). Recent whole-genome studies have confirmed a positive correlation between early replication and the probability of gene expression (3, 4), but have also identified many transcriptionally active and silent genes replicating at all times during S phase. The extent to which replication timing is regulated during development has been even more difficult to substantiate. Very few genes have been demonstrated to replicate at different times in different cell lines, and most genes analyzed do not change replication timing in different cell types (5). Most of these genes have been analyzed in established, often karyotypically unstable cell lines, so the extent to which these differences in replication timing could have resulted from chromosome rearrangements is unclear. Moreover, no examples of programmed changes in replication timing have been observed in a cultured-cell-differentiation system other than those that accompany X-chromosome inactivation (6, 7). These limitations have precluded the ability to directly demonstrate a relationship between replication timing and developmentally regulated patterns of gene expression; thus, it has not been possible to predict how many or what types of genes might be subject to replication timing control. Nonetheless, because chromatin proteins are reassembled at each round of replication, it is reasonable to presume that DNA replication plays some role in epigenetic regulation of gene expression (8).

Addressing the significance of replication timing to developmentally regulated programs of gene expression will require cultured-cell-differentiation systems in which changes in replication control can be elicited in homogeneous cell populations. The recent development of systems for directed changes in cell

fate (9, 10) prompted us to search for autosomal genes that are subject to replication timing changes during the differentiation of mouse embryonic stem cells (ESCs) to neural precursors (neural stem cells, NSCs). Here, we evaluated the replication timing of 54 tissue-specific genes in mouse ESCs before and after differentiation to neural precursors. This analysis identified four genes that switch from early to late replication upon transcriptional down-regulation and five genes that switch from late to early replication upon activation, providing direct evidence for differentiation-induced changes in autosomal replication timing. Intriguingly, differentiation-induced replication timing changes were restricted to isochores rich in AT and long interspersed nuclear elements (LINEs), suggesting a relationship between replication timing and the evolution of metazoan genomes.

Materials and Methods

Cell Culture, Neural Differentiation, and BrdUrd Labeling. ESCs (46C) were routinely grown without feeders on a 0.1% gelatin-coated flask in a leukemia inhibitory factor-supplemented Glasgow minimum essential medium. Differentiation in N2B27 is described in ref. 9. For BrdUrd labeling, both undifferentiated and differentiated cells were incubated with 50 μ M BrdUrd for 1 h before trypsinization and ethanol fixation.

Cell Cycle Fractionation and Isolation of BrdUrd-Labeled DNA. BrdUrd-labeled, fixed cells were resuspended in PBS (2.0 \times 10⁶ cells per ml), stained with propidium iodide (50 μ g/ml) for 30 min in the presence of RNaseA (0.5 mg/ml) and then sorted into two cell cycle fractions (early and late S phase) (Fig. 1A and B) by flow cytometry, as described in refs. 11 and 12. Isolation of BrdUrd-labeled DNA has been described (13, 14). Briefly, 40,000 cells were lysed in SDS-PK buffer (1 M NaCl/10 mM EDTA/50 mM Tris-HCl, pH 8.0/0.5% SDS/0.2 mg/ml PK/50 μ g/ml glycogen), and DNA was extracted by phenol/chloroform extraction followed by ethanol precipitation. Extracted DNA was sonicated, incubated for 20 min at room temperature with anti-BrdUrd antibody (BD Biosciences) in 1 \times immunoprecipitation buffer (10 mM sodium phosphate, pH 7.0/0.14 M NaCl/0.05% Triton X-100), then added to 35 μ g of rabbit anti-mouse IgG (Sigma) for another 20-min incubation. DNA-protein complex was precipitated by centrifugation, washed once with 1 \times immunoprecipitation buffer, and resuspended in digestion buffer (50 mM Tris-HCl, pH 8.0/10 mM EDTA/0.5% SDS/0.25 mg/ml PK) for overnight protein digestion at 37°C. Finally, immunoprecipitated DNA was collected by ethanol precipitation and resuspended in Tris-EDTA at a concentration of 250 cell equivalents per μ l.

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: LINE, long interspersed nuclear element; GCL, GC-low; GCH, GC-high; NSC, neural stem cell; ESC, embryonic stem cell; txStart, transcription start position.

*To whom correspondence should be addressed. E-mail: gilbertd@upstate.edu.

© 2004 by The National Academy of Sciences of the USA




Fig. 1. Experimental scheme of replication-timing analysis before and after differentiation. (A and B) Undifferentiated (A) and neural differentiated (B) ESCs were pulse-labeled with BrdUrd and then separated into populations from early or late S phase by flow cytometry. The white dotted lines represent the sorting windows for fractions. (C) Differentiation state as determined by RT-PCR. *Oct4* (ESC-specific) is down-regulated after differentiation, whereas neural precursor-specific *Sox1* is up-regulated. Samples marked –RT did not contain reverse transcriptase in the reaction. (D) Replication-timing analysis of control genes, α -globin (early-replicating in ESC), β -globin (late-replicating in ESC), and mitochondrial DNA (mito; replicates throughout the entire cell cycle) (28). The average and SEM of early S phase abundance (percentage of the total) were measured as described in *Materials and Methods*. Undiff., undifferentiated ESC; diff., differentiated ESC; E, early S-phase fraction; L, late S-phase fraction; PI, propidium iodide.

Replication-Timing Analysis by PCR. A control PCR (35 cycles) experiment was performed to confirm the enrichment of α -globin, β -globin, and mitochondrial DNA sequences in the expected fractions of immunoprecipitated BrdUrd-labeled DNA samples, early S phase, late S phase, and both fractions, respectively. We then determined whether a gene replicated in the first half or the second half of S phase based on conventional PCRs (35–42 cycles, determined empirically for each primer set) and subsequent ethidium bromide gel staining. For α -globin, β -globin, and mitochondrial DNA, a 23-cycle PCR followed by Southern hybridization (5) gave results comparable to ethidium bromide gel staining (data not shown). To quantify the relative enrichment in each fraction, we measured the intensity of the PCR products by using the image processing software IMAGEJ (<http://rsb.info.nih.gov/ij>). The relative abundance was determined by calculating the early S-phase percentage of total: (intensity of early S phase)/(intensities of early and late S phase combined). A percentage $>60\%$ and $<40\%$ in early S fraction was defined as early replicating (E) and late replicating (L), respectively. We categorized genes that showed 40–60% enrichment in early S phase as early replicating (E) because cells are synchronized after the completion of a 1-h BrdUrd pulse-labeling and genes that replicated in early middle S phase move into late S phase by the end of the pulse-labeling period.

Gene Expression Analysis by RT-PCR. RNA isolation was done by using a RNeasy RNA extraction kit (Qiagen, Valencia, CA). Extracted RNA was treated with DNaseI to eliminate genomic DNA, extracted with phenol/chloroform, and ethanol precipitated. Reverse transcription was performed by using random hexamer (Invitrogen) and SuperScript III (Invitrogen). For every primer set, we initially used serial dilutions of cDNA for

PCR and stopped the reaction at every two or three cycles between 21 and 30 cycles to determine the cycle numbers that fall within the exponential range of the PCR, where the amount of PCR products of 2-fold serial dilutions should stay in a linear range. After cycle number determination, cDNA derived from undifferentiated and differentiated ESCs was analyzed by PCR and quantified by using IMAGEJ as in the replication timing analysis. For the GC-low (GCL) genes and the GC-high (GCH) genes analyzed in this study, a difference of >2.5 - and 1.5-fold was defined as tissue-specific, respectively (GCL genes showed a higher average fold-change than GCH genes).

Gene Selection, GC Content Calculation, and Primer Design. For gene selection, we used three existing microarray databases of genes enriched in mouse ESCs and animal-derived NSCs (15–17) and selected genes that were reproducibly defined as either ESC-specific or NSC-specific in at least two of the three studies. For the ESC-GCL, ESC-GCH, NSC-GCL, and NSC-GCH categories, 36, 32, 24, and 25 primer sets, respectively, were initially designed, and a total of 54 genes presented in this study gave PCR results. For GC content calculations, 400 kb of surrounding DNA [200 kb upstream and downstream of the transcription start position (txStart)] was analyzed, based on the October 2003 mouse genome assembly (build 32) of the University of California, Santa Cruz, genome browser (<http://genome.ucsc.edu>). For high-throughput GC content calculations of large numbers of genes, 20-kb windows containing the txStart and transcription end positions were analyzed. The txStart and transcription end positions can be found on the University of California, Santa Cruz, genome browser. For those genes analyzed out to 400-kb windows, 20-kb windows were found to be representative of the entire isochore (Tables 2 and 3, which are published as supporting information on the PNAS web site). Primers were designed to amplify 200- to 650-bp fragments by PRIMER3 (<http://frodo.wi.mit.edu/cgi-bin/primer3/primer3-www.cgi>).

Analysis of Point Mutation Rate, LINE Density, and Gene Frequency Versus GC Content. A data set containing normalized 4-fold substitution rates of 14,790 mouse/human orthologous genes was kindly provided by H. Li and J. Chuang (University of California, San Francisco) (18). We were able to identify the GC content of a 20-kb window surrounding the txStart position for 8,097 genes, and these genes were used for the analysis. For LINE density, chrN_rmsk.txt files were downloaded from the University of California, Santa Cruz, genome browser (build 32, October 2003), and the percentage of repetitive sequences per 400 kb surrounding the txStart position was calculated for each gene and plotted against average GC content of the surrounding 400-kb region. For gene frequency versus GC content, 16,652 National Center for Biotechnology Information Reference Sequence (RefSeq) genes and GC content value of the 20-kb windows surrounding the txStart position were used.

Results and Discussion

Predicting Genes That Switch Replication Timing. Because there are so few examples of genes that change their replication timing, one challenge was how to predict which genes, if any, might be subject to developmental control over DNA replication in any given differentiation system. We reasoned that clues might be derived from a comparison of the α -globin and β -globin genes. These genes encode subunits of the same protein complex and are coordinately regulated, yet their replication timing regulation is very different: α -Globin is early-replicating and euchromatic in all tissues, whereas β -globin is late-replicating and heterochromatic in most tissues and early-replicating in erythroid cells (19–22). This example renders unlikely any hypotheses that relate replication-timing switches to gene function or expression pattern. One notable difference between these genes




Fig. 3. Replication-timing analysis of NSC- and ESC-specific AT-rich (GCL) genes analyzed as for Fig. 2. E, early-replicating; L, late-replicating.

replication timing during development. Genes that remain early-replicating after down-regulation may remain poised for expression or may be adjacent to genes that remain expressed. We have only examined one stage of differentiation, and these genes may switch to late replication in other tissues where they are repressed. Moreover, two of these genes (*Trap1a* and *Car8*) displayed changes in replication timing that were greater than those observed for most GCH genes, but, because they were confined to early S phase, these genes were categorized as early in both differentiation states for the reasons stated above. Among those GCL genes that were expressed in both undifferentiated and differentiated ESCs, all but one remained early-replicating (Fig. 4), and the mean difference in replication timing between differentiation states for these genes was significantly smaller than for GCL genes that did change expression (7 ± 2 versus 23 ± 3). These observations strongly suggest that genes residing within AT-rich isochores have a high propensity to switch replication timing coincident with changes in transcriptional state.

Our analysis revealed five exceptional genes that were late-replicating and expressed (Fig. 3: *Ptgs2*, *Nr0b1*, *Sprr2a*, and *A2m*; Fig. 4: *C1qr1*). No such genes were found among the GCH group. Importantly, robust expression of late-replicating genes was primarily observed in ESCs (four of five genes) (Fig. 3: *Nr0b1*, *Sprr2a*, and *A2m*; Fig. 4: *C1qr1*). Because examples of late-replicating, expressed genes have been rare in the literature (Table 2), our observation raises the intriguing possibility that late-replicating regions are not as readily assembled into silent chromatin in pluripotent ESCs (31). Whole-genome analyses of




Fig. 4. Analysis of AT-rich genes that are expressed before and after differentiation (Both-GCL) as for Fig. 2. E, early-replicating; L, late-replicating.

replication timing versus gene expression in ESCs and NSCs should assess this possibility by determining whether the positive correlation identified in established *Drosophila* (3) and human (4) cell lines is less prominent in ESCs compared with NSCs.

Our results are summarized in Table 1. Strikingly, for genes that changed their transcriptional state, replication-timing changes were observed with 10 of 19 GCL genes and none of the 18 GCH genes (Fisher's exact test, $P < 0.001$). Because early replication is presumed to set up a permissive chromatin structure but is not sufficient for gene expression and some of these loci may be adjacent to expressed genes, a certain number of early-replicating nonexpressed GCL genes are expected. These results indicate that differentiation-induced changes in replication control are limited to genes that reside within AT-rich isochores.

A Threshold LINE Density for Replication-Timing Switches. Are there predictive characteristics other than GC content that define the class of genes subject to developmentally regulated replication-timing switches? In the course of these studies, we noted that our GCL genes were much larger with more intronic sequences (Table 3). We therefore used genome sequencing data to examine the complete set of mouse GCH and GCL genes for

Table 1. Summary

Gene category	Replication timing				Switches observed
	L→E	L→L	E→L	E→E	
ESC-GCL	1	2	4	4	
NSC-GCL	5	1	0	2	10/19 (GCL)
ESC-GCH	0	0	0	9	
NSC-GCH	0	0	0	9	0/18 (GCH)

Tissue-specific genes analyzed in this study were categorized according to their gene expression pattern and GC content, and their replication timing results are shown. Replication-timing switches were only observed within the GCL gene category ($P < 0.001$, Fisher's exact test). E and L represent early- and late-replicating, respectively. For instance, L→E indicates switch from late replication to early replication upon differentiation.

determine whether this relationship bears out in other vertebrates that have different isochore structure and other types of transposons that may have different sequence preferences (46).

In summary, our results provide the first demonstration of replication timing changes induced by differentiation in culture, setting the stage for mechanistic studies of the relationship between replication timing and developmentally regulated patterns of gene expression. In doing so, we provide direct evidence for the longstanding notion that replication timing changes are a significant part of mammalian development, as more than one-third of all genes are potential candidates for such changes (Fig. 5D). Moreover, we also offer a means to predict which genes are subject to this type of regulation in any given mammalian differentiation system. Finally, this class of genes is rich

in LINEs, and their structure implies that they have been subjected to rapidly evolving changes in developmentally regulated gene-expression patterns, a driving force in metazoan evolution.

We thank A. Smith and M. Stavridis (University of Edinburgh, Edinburgh) for the 46C cell line and technical advice; H. Li and J. Chuang for providing mutational analysis data; Z. Lai and N. Gonchoroff for generously assisting with flow cytometry; S. Fiering and J. Goodier for helpful discussion; B. Knox, T. Sasaki, A. Fisher, and V. Azuara for comments on the manuscript; and J. Moran, H. Li, and J. Chuang for discussion and comments on the manuscript. This work was supported by National Institutes of Health Grant GM-57233-01 (to D.M.G.) and by a predoctoral fellowship from the Japan Society for the Promotion of Science (to I.H.).

1. Gilbert, D. M. (2002) *Curr. Opin. Cell Biol.* **14**, 377–383.
2. Goren, A. & Cedar, H. (2003) *Nat. Rev. Mol. Cell Biol.* **4**, 25–32.
3. Schubeler, D., Scalzo, D., Kooperberg, C., van Steensel, B., Delrow, J. & Groudine, M. (2002) *Nat. Genet.* **32**, 438–442.
4. Woodfine, K., Fiegler, H., Beare, D. M., Collins, J. E., McCann, O. T., Young, B. D., Debernardi, S., Mott, R., Dunham, I. & Carter, N. P. (2004) *Hum. Mol. Genet.* **13**, 191–202.
5. Azuara, V., Brown, K. E., Williams, R. R., Webb, N., Dillon, N., Festenstein, R., Buckle, V., Merckenschlager, M. & Fisher, A. G. (2003) *Nat. Cell Biol.* **5**, 668–674.
6. Keohane, A. M., O'Neill, L. P., Belyaev, N. D., Lavender, J. S. & Turner, B. M. (1996) *Dev. Biol.* **180**, 618–630.
7. Brockdorff, N. (2002) *Trends Genet.* **18**, 352–358.
8. McNairn, A. J. & Gilbert, D. M. (2003) *BioEssays* **25**, 647–656.
9. Ying, Q. L., Stavridis, M., Griffiths, D., Li, M. & Smith, A. (2003) *Nat. Biotechnol.* **21**, 183–186.
10. Rathjen, J., Haines, B. P., Hudson, K. M., Nesci, A., Dunn, S. & Rathjen, P. D. (2002) *Development (Cambridge, U.K.)* **129**, 2649–2661.
11. Gilbert, D. M. (1986) *Proc. Natl. Acad. Sci. USA* **83**, 2924–2928.
12. Gilbert, D. & Cohen, S. (1987) *Cell* **50**, 59–68.
13. Cimbora, D. M., Schubeler, D., Reik, A., Hamilton, J., Francastel, C., Epner, E. M. & Groudine, M. (2000) *Mol. Cell. Biol.* **20**, 5581–5591.
14. Hansen, R., Canfield, T., Lamb, M., Gartler, S. & Laird, C. (1993) *Cell* **73**, 1403–1409.
15. Ramalho-Santos, M., Yoon, S., Matsuzaki, Y., Mulligan, R. C. & Melton, D. A. (2002) *Science* **298**, 597–600.
16. Ivanova, N. B., Dimos, J. T., Schaniel, C., Hackney, J. A., Moore, K. A. & Lemischka, I. R. (2002) *Science* **298**, 601–604.
17. Fortunel, N. O., Otu, H. H., Ng, H. H., Chen, J., Mu, X., Chevassut, T., Li, X., Joseph, M., Bailey, C., Hatzfeld, J. A., et al. (2003) *Science* **302**, 393.
18. Chuang, J. H. & Li, H. (2004) *PLoS Biol.* **2**, E29.
19. Furst, A., Brown, E. H., Braunstein, J. D. & Schildkraut, C. L. (1981) *Proc. Natl. Acad. Sci. USA* **78**, 1023–1027.
20. Dhar, V., Mager, D., Iqbal, A. & Schildkraut, C. L. (1988) *Mol. Cell. Biol.* **8**, 4958–4965.
21. Forrester, W. C., Epner, E., Driscoll, M. C., Enver, T., Brice, M., Papayanopoulou, T. & Groudine, M. (1990) *Genes Dev.* **4**, 1637–1649.
22. Brown, K. E., Amoils, S., Horn, J. M., Buckle, V. J., Higgs, D. R., Merckenschlager, M. & Fisher, A. G. (2001) *Nat. Cell Biol.* **3**, 602–606.
23. Hatton, K. S., Dhar, V., Brown, E. H., Iqbal, M. A., Stuart, S., Didamo, V. T. & Schildkraut, C. L. (1988) *Mol. Cell. Biol.* **8**, 2149–2158.
24. Selig, S., Okumura, K., Ward, D. C. & Cedar, H. (1992) *EMBO J.* **11**, 1217–1225.
25. Watanabe, Y., Fujiyama, A., Ichiba, Y., Hattori, M., Yada, T., Sakaki, Y. & Ikemura, T. (2002) *Hum. Mol. Genet.* **11**, 13–21.
26. Eyre-Walker, A. (1992) *Nucleic Acids Res.* **20**, 1497–1501.
27. Eyre-Walker, A. & Hurst, L. D. (2001) *Nat. Rev. Genet.* **2**, 549–555.
28. Aladjem, M. I., Rodewald, L. W., Lin, C. M., Bowman, S., Cimbora, D. M., Brody, L. L., Epner, E. M., Groudine, M. & Wahl, G. M. (2002) *Mol. Cell. Biol.* **22**, 442–452.
29. Panning, M. M. & Gilbert, D. M. (2005) *J. Cell. Biochem.*, in press.
30. Dimitrova, D. S. & Gilbert, D. M. (1999) *Mol. Cell* **4**, 983–993.
31. Gasser, S. M. (2002) *Science* **296**, 1412–1416.
32. Eisenberg, E. & Levanon, E. Y. (2003) *Trends Genet.* **19**, 362–365.
33. Shapiro, M. D., Marks, M. E., Peichel, C. L., Blackman, B. K., Nereng, K. S., Jonsson, B., Schluter, D. & Kingsley, D. M. (2004) *Nature* **428**, 717–723.
34. Shubin, N. H. & Dahn, R. D. (2004) *Nature* **428**, 703–704.
35. Moran, J. & Gilbert, N. (2002) in *Mobile DNA II*, eds. Craig, N. L., Craigie, R., Gellert, M. & Lambowitz, A. (Am. Soc. Microbiol. Press, Washington, D.C.), pp. 836–869.
36. Han, J. S., Szak, S. T. & Boeke, J. D. (2004) *Nature* **429**, 268–274.
37. Waterston, R. H., Lindblad-Toh, K., Birney, E., Rogers, J., Abril, J. F., Agarwal, P., Agarwala, R., Ainscough, R., Alexandersson, M., An, P., et al. (2002) *Nature* **420**, 520–562.
38. Allen, E., Horvath, S., Tong, F., Kraft, P., Spiteri, E., Riggs, A. D. & Marahrens, Y. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 9940–9945.
39. Lynch, M. & Conery, J. S. (2003) *Science* **302**, 1401–1404.
40. Feng, Q., Moran, J. V., Kazazian, H. H., Jr., & Boeke, J. D. (1996) *Cell* **87**, 905–916.
41. Schramke, V. & Allshire, R. (2004) *Curr. Opin. Genet. Dev.* **14**, 174–180.
42. Xie, Z., Johansen, L. K., Gustafson, A. M., Kasschau, K. D., Lellis, A. D., Zilberman, D., Jacobsen, S. E. & Carrington, J. C. (2004) *PLoS Biol.* **2**, E104.
43. Wakimoto, B. T. (1998) *Cell* **93**, 321–324.
44. Smit, A. F., Toth, G., Riggs, A. D. & Jurka, J. (1995) *J. Mol. Biol.* **246**, 401–417.
45. Bailey, J. A., Carrel, L., Chakravarti, A. & Eichler, E. E. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 6634–6639.
46. Jaillon, O., Aury, J. M., Brunet, F., Petit, J. L., Stange-Thomann, N., Mauceli, E., Bouneau, L., Fischer, C., Ozouf-Costaz, C., Bernot, A., et al. (2004) *Nature* **431**, 946–957.

Table 2. GC content of mammalian genes whose replication timing has been analyzed in different cell lines

Gene	Gene abbr.	Chr	Replication timing switch	GC% 400 kb	GC% txStart	GC% txEnd	%LINE	%Lx	Average LINE length (bp)	Ref.
AFP	Afp	chr5	yes	41.2	41.6	41.7	14.4%	4.4%	429	1
albumin	Alb1	chr5	yes	41.2	39.3	39.3	13.8%	4.1%	427	1
β-globin	Hbb-b1	chr7	yes	38.3	40.9	40.9	31.1%	12.5%	569	1
TCR beta chain, constant region (Tcβ)	X01643	chr6	yes	39.3	39.0	44.0	26.9%	8.8%	498	1
Vβ19	U59155	chr6	yes	40.6	39.2	39.2	49.8%	17.1%	772	1
Immunoglobulin heavy chain, VHT15	AF118974	chr12	yes	38.9	39.9	N.D.	40.6%	17.3%	636	1
DHFR	Dhfr	chr13	yes	42.3	43.4	43.1	15.3%	2.7%	461	1
CFTR	Cfr	chr6	yes	38.3	39.5	38.9	24.0%	6.7%	535	2
MUP	Mup1	chr4	no (late)	40.7	40.0	39.3	33.7%	0.0%	1184	1
Amylase	Amy1	chr3	no (late)	37.4	37.8	34.8	39.9%	14.5%	752	1
Ig Kappa, constant region (Cκ-Jκ)	U17383	chr6	no (early)	43.3	39.8	39.8	9.4%	2.4%	350	1
CD45	L36091	chr1	no (early)	38.8	36.3	37.7	14.2%	5.4%	510	3
Rag-1	Rag1	chr2	no (early)	43.0	42.1	42.0	11.7%	3.2%	417	3
CD8β	BC030679	chr6	no (early)	43.2	44.9	43.4	7.5%	1.9%	267	3
β5 locus	Vpreb1	chr16	no (early)	42.3	45.8	45.8	27.4%	6.3%	851	3
TdT	Dntt	chr19	no (early)	44.8	45.8	45.8	4.8%	1.5%	255	3
β-globin	Hba-a1	chr11	no (early)	46.7	47.2	47.2	5.1%	0.7%	242	1
Immunoglobulin, Cλ	AB058113	chr12	no (early)	49.5	50.3	50.3	11.2%	3.7%	608	1
C2	C2	chr17	no (early)	50.4	47.9	51.1	1.4%	0.5%	228	1
C4	C4	chr17	no (early)	50.9	53.4	53.4	2.2%	0.9%	232	1
Thy-1	Thy1	chr9	no (early)	47.7	49.7	49.7	4.0%	1.1%	249	1
Tyrosine amino-transferase	Tat	chr8	no (early)	45.1	45.6	45.6	6.0%	1.2%	333	1
Metallothionein	Mt1	chr8	no (early)	45.5	48.9	48.9	5.4%	1.1%	230	1
CD19	Cd19	chr7	no (early)	46.7	48.7	46.7	5.9%	3.3%	314	3
CD4	Cd4	chr6	no (early)	48.7	53.7	46.6	3.0%	1.4%	208	3
α1-antitrypsin	M25529	chr12	no (late)	44.3	48.6	48.6	12.4%	0.8%	520	1

From this sampling, every gene found to change replication timing was AT-rich (<43% GC, 8/8). For Vβ19, Ig heavy chain, CFTR, Amylase, Ig Kappa, CD8β, CD19, and α1-antitrypsin, genomic sequence within the 400 kb region surrounding the gene was partially incomplete. These incomplete sequences were excluded from the GC% and LINE density analysis. Gene abbr., gene name abbreviation or accession numbers used in the UCSC genome database; Chr, chromosome; “No (early)” and “no (late),” either early or late replicating in all cell lines examined, respectively; GC% 400 kb, GC% of the 400 kb region surrounding the transcription start position (200 kb upstream and downstream); GC% txStart and GC% txEnd, GC% of the 20

kb region surrounding the transcription start and end sites, respectively; %LINE and %Lx, the density of LINE and Lx sequence in base pairs, respectively, in the 400 kb region; Average LINE length (bp), average length of LINEs within the 400 kb region; N.D., not determined.

1. Hatton, K. S., Dhar, V., Brown, E. H., Iqbal, M. A., Stuart, S., Didamo, V. T. & Schildkraut, C. L. (1988) *Mol. Cell. Biol.* **8**, 2149-2158.
2. Selig, S., Okumura, K., Ward, D. C. & Cedar, H. (1992) *EMBO J.* **11**, 1217-1225.
3. Azuara, V., Brown, K. E., Williams, R. R., Webb, N., Dillon, N., Festenstein, R., Buckle, V., Merkenschlager, M. & Fisher, A. G. (2003) *Nat. Cell Biol.* **5**, 668-674.

Table 3. Sequence characteristics of genes analyzed in this study

Gene category	Gene name	Chr	Replication timing	GC% 400 kb	GC% txStart	GC% txEnd	tx length (bp)	Intron length (bp)	Intron#	%LINE	%Lx	Average LINE length (bp)
ESC-GCH	Pem	chrX random	E -> E	N.D.	47.5	47.5	4031	3281	3	N.D.	N.D.	N.D.
ESC-GCH	Ebaf	chr1	E -> E	46.0	50.2	50.2	3379	1759	3	4.4%	0.8%	311
ESC-GCH	Dmrt1	chr19 random	E -> E	N.D.	50.4	50.4	4139	3478	1	N.D.	N.D.	N.D.
ESC-GCH	Tdgf1/Cripto	chr9	E -> E	47.3	47.3	47.3	6465	4624	5	5.1%	0.7%	260
ESC-GCH	Tcfap2c	chr2	E -> E	49.4	52.1	52.1	8996	6192	6	1.7%	0.1%	224
ESC-GCH	2010004A03Rik	chr8	E -> E	47.3	51.8	48.3	64036	62240	7	5.0%	0.3%	247
ESC-GCH	Utf1	chr7	E -> E	48.7	50.3	50.3	1257	97	1	4.0%	1.1%	307
ESC-GCH	Ptgis	chr2	E -> E	49.5	53	48.8	37342	35591	9	2.9%	0.6%	195
ESC-GCH	Epha2	chr4	E -> E	50.4	53.8	54.2	28139	24232	16	1.9%	0.5%	146
Both-GCH	Pcolce	chr5	E -> E	48.3	49.7	49.7	6251	4759	8	4.0%	1.8%	266
Both-GCH	Ltbr	chr6	E -> E	48.8	50.3	50.3	7360	5289	9	3.9%	1.2%	194
Both-GCH	Ddr1	chr17	E -> E	48.6	50.8	50.2	23318	19689	17	2.3%	1.3%	223
Both-GCH	Dll1	chr17	E -> E	44.8	49.8	49.8	7900	5043	10	6.8%	2.0%	326
NSC-GCH	Mmp15	chr8	E -> E	47.7	51.9	51.5	20726	17982	10	4.6%	1.4%	222
NSC-GCH	Slc25a10	chr11	E -> E	50.3	51.7	51.7	7294	5325	10	1.3%	0.3%	197
NSC-GCH	Nfatc1	chr18	E -> E	46.8	51.1	52.3	60312	56234	7	6.5%	1.3%	366
NSC-GCH	Cdh4	chr2	E -> E	48.0	50.7	50.1	453715	450431	15	4.0%	1.3%	291
NSC-GCH	Bmp1	chr14	E -> E	47.9	52.1	50.5	44923	41957	19	5.6%	1.9%	317
NSC-GCH	Fgfr1	chr8	E -> E	45.5	49.1	50.1	41934	39408	16	3.6%	1.1%	201
NSC-GCH	Gpr56	chr8	E -> E	49.4	49.4	48.9	10942	7736	11	3.5%	0.5%	217
NSC-GCH	Lmo1	chr7	E -> E	44.2	48.7	48.7	2139	1605	1	18.4%	4.2%	549
NSC-GCH	Cspg3	chr8	E -> E	49.8	51.5	51.2	25845	20596	14	0.7%	0.0%	118
NSC-GCL	Ptn	chr6	L -> E	41.3	39.4	41.5	95821	93800	4	14.7%	4.3%	446
NSC-GCL	A1426782/Tm4sf12	chr6	L -> E	39.0	39.2	44.1	81120	78630	8	19.2%	9.3%	541
NSC-GCL	Postn	chr3	L -> E	40.5	40.8	40.3	29926	26724	21	13.6%	6.6%	384
NSC-GCL	Ptprz1	chr6	L -> E	39.4	39.9	37.8	119582	3724	10	26.4%	7.8%	513
NSC-GCL	Lox	chr18	L -> E	44.1	36.8	44.3	13637	8867	6	14.6%	3.8%	504
NSC-GCL	Mtap2	chr1	E -> E	38.8	38.2	39.1	113500	107617	14	13.9%	5.7%	474
NSC-GCL	Sox6	chr7	E -> E	42.5	42.4	42.3	52133	51286	5	6.9%	1.7%	440
NSC-GCL	Ptgs2	chr1	L -> L	40.0	39.1	39.1	7887	3744	9	24.3%	7.2%	639
ESC-GCL	Rex1/Zfp42	chr8	E -> L	41.6	40.5	44.9	11890	10145	4	17.6%	6.2%	451
ESC-GCL	Rex2	chr4	E -> L	40.4	39.8	38.3	85827	84760	5	15.9%	10.0%	511
ESC-GCL	Crisp1	chr17	E -> L	37.3	35.9	35.9	25440	24043	7	35.8%	8.5%	705
ESC-GCL	Adcy8	chr15	E -> L	41.1	41.8	43.3	218011	213809	15	14.5%	3.8%	436
ESC-GCL	Trap1a	chrX	E -> E	39.6	41.5	41.5	4483	3309	2	31.0%	10.9%	498
ESC-GCL	Spp1	chr5	E -> E	40.9	40.7	43.4	4852	3345	5	24.9%	7.4%	727
ESC-GCL	Car8	chr4	E -> E	40.8	41.9	41.7	93503	91824	8	24.9%	9.0%	560
ESC-GCL	Gbp3	chr3	E -> E	41.3	41.9	41.5	13057	10392	12	19.6%	8.0%	584
ESC-GCL	Nr0b1	chrX	L -> L	39.6	40.1	40.1	4160	2376	1	31.3%	15.4%	469
ESC-GCL	Sprr2a	chr3	L -> L	39.5	39.2	39.2	3957	3113	1	26.9%	13.1%	438
ESC-GCL	A2m	chr6	L -> E	42.4	41.1	40.7	39194	34589	35	24.4%	4.6%	724
Both-GCL	1200008D14Rik	chr16	E -> E	39.8	42.7	40.8	59374	56483	12	37.0%	8.2%	796
Both-GCL	1700034H14Rik	chr18	E -> E	42.2	42.2	42.2	3128	2553	5	28.6%	5.1%	674
Both-GCL	AA543502	chr8	E -> E	41.1	39.4	39.4	N.D.	N.D.	N.D.	19.1%	4.4%	591
Both-GCL	Ctgf	chr10	E -> E	41.3	43.5	41.3	3174	845	5	12.5%	3.2%	468
Both-GCL	Ddx4	chr13	E -> E	42.6	40.1	42.8	53966	51207	20	10.2%	3.0%	356
Both-GCL	Rbpms	chr8	E -> E	44.6	42.8	N.D.	145429	144330	8	2.9%	0.6%	231
Both-GCL	Tfpi	chr2	E -> E	39.7	36.1	41.8	41315	39823	8	14.3%	7.3%	425
Both-GCL	X98848	chrX	E -> E	41.4	40.2	46.3	4194	3724	3	49.9%	15.5%	653
Both-GCL	Mme	chr3	E -> E	38.9	39.1	36.7	86360	82120	22	25.6%	3.7%	652
Both-GCL	Egfr	chr11	E -> E	40.8	42.6	41.1	135372	132778	15	22.2%	7.4%	522
ESC-GCL	Col1a2	chr6	E -> E	40.3	40.4	40.4	34972	30706	51	13.9%	3.0%	509
Both-GCL	Terf1	chr1	E -> E	41.2	41.6	40.6	37773	36141	9	17.2%	4.0%	662
Both-GCL	C1qr1	chr2	L -> L	42.9	42.8	42.8	6893	3823	2	13.7%	4.8%	456

	tx Length (bp)	Intron Length (bp)	Intron#	%LINE	%Lx	Average LINE Length (bp)
GCH genes analyzed	39566	37161	9.0	4.3%	1.1%	259
GCL genes analyzed	52578	46472	10.7	20.8%	6.6%	533
RefSeq GCH	24487	22194	8.4	N.D.	N.D.	N.D.
RefSeq GCL	53383	51200	6.9	N.D.	N.D.	N.D.

For *Tcfap2c*, *Pcolce*, *Ddr1*, *Nfatc1*, *Lmo1*, *Lox*, *Rex2*, *Crisp1*, *Adcy8*, and *Rbpms*, genomic sequence within the 400 kb region surrounding the gene was partially incomplete. These incomplete sequences were excluded from the GC% and LINE density analysis. The bottom table shows average values of 22 GCH genes and 32 GCL genes analyzed in this study, as well as those of 5533 GCL and 6003 GCH RefSeq genes. Chr, chromosome; GC% 400 kb, average GC% of the 400 kb region surrounding the transcription start position (200 kb upstream and downstream); GC% txStart and GC% txEnd, GC% of the 20 kb region surrounding the transcription start and end sites, respectively; tx length (bp), length of the transcript; Intron#, number of introns in the transcript; %LINE and %Lx, the density of LINE and Lx sequence in base pairs, respectively, in the 400 kb region; Average LINE length (bp), average length of LINES within the 400 kb region; N.D., not determined.

Fig. 6. Sequence characteristics of genes within GCL (GC-Low) isochores. The mutation rates (normalized substitution rates at 4-fold sites) of 8097 genes (1) plotted against their GC% of the 20-kb region surrounding the transcription start (txStart) position, which is defined by the UCSC genome browser. No correlation was observed ($R^2 = 0.0523$).

1. Chuang, J. H. & Li, H. (2004) *PLoS Biol.* **2**, E29.

